

CAUTION

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING



Affirm[®]
Insecticide

syngenta.

ACTIVE CONSTITUENT:

17 g/L EMAMECTIN present as EMAMECTIN BENZOATE

GROUP **6** INSECTICIDE

For the control of *Heliothis* in cotton and the suppression of Mites and Green Mirids in Cotton

Syngenta Crop Protection Pty Limited
Level 1, 2-4 Lyonpark Road, Macquarie Park NSW 2113

In a transport emergency dial 000, Police or Fire Brigade
For specialist advice in an emergency only, call 1800 033 111 (24 hours)

APVMA Approval No.: 51321/20/1005

TM

DIRECTIONS FOR USE

Restraint: DO NOT apply to under visible stress

Restraint: DO NOT apply to cotton under visible stress.

Crop	Pest	Rate	Critical Comments
Cotton Qld, NSW, WA, NT only	Heliothis <i>Helicoverpa</i> <i>armigera</i> , <i>Helicoverpa</i> <i>punctigera</i>	550 mL/ha or 700 mL/ha Add a non-ionic surfactant at recommended label rate	Timing is important. For best results apply at or just prior to the anticipated time of hatching of <i>Helicoverpa</i> eggs as indicated by egg levels or scouting. As earlier larval stages are easier to control, DO NOT target larvae larger than 5 mm (very small to small). Large larvae and larvae feeding within bolls and squares may not be controlled. Larvae must feed on treated areas in order to achieve effective control. Where high egg lays occur the addition of an ovicide is recommended. Use the lower rate on light infestations. Make no more than 4 applications per season. AFFIRM should be used according to the cotton Best Management Practices Manual.
	Green Mirids (<i>Creontiades</i> <i>dilutus</i>) Suppression only		Apply to developing Green Mirid populations that are predominantly nymphs. Maximum effect is not achieved until 5 to 7 days after application. Use the higher rate on heavier populations.
	Mites Suppression only		Applications of AFFIRM as outlined above for <i>Heliothis</i> will reduce the rate of mite development in treated cotton.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION

WITHHOLDING PERIODS

DO NOT HARVEST, GRAZE OR CUT FOR STOCK FOOD FOR 4 WEEKS AFTER APPLICATION

DO NOT FEED COTTON TRASH FROM TREATED AREAS TO ANIMALS, INCLUDING POULTRY

GENERAL INSTRUCTIONS

Mixing

Add the required amount of AFFIRM to a partly filled spray tank, and then add the remainder of the water.

Application

To be effective AFFIRM requires thorough spray coverage. Ensure that equipment is properly calibrated to give an even distribution at the correct volume.

Ground Application: Apply in at least 80 L spray/ha

Air Application: Apply in at least 20 L spray/ha

Compatibility

AFFIRM is compatible with Ovasyn* EC. As formulations of other manufacturer's products are beyond the control of Syngenta, and the quality of water may vary with location, all mixtures should be tested prior to mixing commercial quantities.

Insecticide Resistance Warning

GROUP	6	INSECTICIDE
-------	----------	-------------

For insecticide resistance management AFFIRM INSECTICIDE is a Group 6 insecticide. Some naturally occurring insect biotypes resistant to AFFIRM and other Group 6 insecticides may exist through normal genetic variability in any insect population. The resistant individuals can eventually dominate the insect population if AFFIRM and other Group 6 insecticides are used repeatedly. The effectiveness of AFFIRM on resistant individuals could be significantly reduced. Since occurrence of resistant individuals is difficult to detect prior to use Syngenta Crop Protection Pty Limited accepts no liability for any losses that may result from failure of AFFIRM to control resistant insects. AFFIRM may be subject to specific resistance management strategies. For further information contact your supplier, Syngenta Crop Protection representative or local agricultural department agronomist.

PRECAUTION

Human flagging is not supported unless flaggers are protected by engineering controls such as vehicles with cabs.

Re-entry Period

DO NOT allow entry into treated areas for 12 hours after treatment. When prior entry is necessary, wear cotton overalls buttoned to the neck and wrist and chemical resistant gloves.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

Dangerous to fish and other aquatic organisms. DO NOT contaminate streams, rivers or waterways with the chemical or used containers. DO NOT apply under weather conditions or from spraying equipment that may cause spray to drift from the target area. A spray drift minimisation strategy should be employed at all times when aerially applying sprays to, or near, sensitive areas. The strategy envisaged is exemplified by the cotton industry's Best Management Practice Manual.

PROTECTION OF LIVESTOCK

Dangerous to bees. DO NOT spray any plants in flower while bees are foraging.

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well ventilated area. DO NOT store for prolonged periods in direct sunlight.

Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

SAFETY DIRECTIONS

Will damage the eyes. Will irritate the skin. Avoid contact with eyes and skin. When opening the container, and preparing spray wear:

- cotton overalls buttoned to the neck and wrist
- washable hat
- elbow length PVC gloves and
- goggles

When using the prepared spray wear:

- cotton overalls buttoned to the neck and wrist
- washable hat
- elbow-length PVC gloves

If product in eyes, wash it out immediately with water. If product on skin, immediately wash area with soap and water. Wash hands after use. After each day's use, wash gloves, goggles and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131 126.

MATERIAL SAFETY DATA SHEET

If additional hazard information is required, refer to the Material Safety Data Sheet. For a copy phone 1800 067 108 or visit our website at www.syngenta.com.au

MANUFACTURER'S WARRANTY AND EXCLUSION OF LIABILITY

Syngenta has no control over storage, handling and manner of use of this product. Where this material is not stored, handled or used correctly and in accordance with directions, no express or implied representations or warranties concerning this product (other than non-excludable statutory warranties) will apply. Syngenta accepts no liability for any loss or damage arising from incorrect storage, handling or use.

Product names marked ® or ™, the ALLIANCE FRAME the SYNGENTA Logo and the PURPOSE ICON are Trademarks of a Syngenta Group Company



*Trademark