

# CAUTION

KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING



# Glyphosate Trimesium 480

**HERBICIDE**

ACTIVE CONSTITUENT: 480 g/L GLYPHOSATE-TRIMESIUM

|       |          |           |
|-------|----------|-----------|
| GROUP | <b>M</b> | HERBICIDE |
|-------|----------|-----------|

*A non-selective foliar applied systemic herbicide for the control of a wide range of annual and perennial weeds as per Directions for Use*


**IMPORTANT:** Read the attached booklet before use

## 20 LITRES

**Syngenta Crop Protection Pty Limited**  
Level 1, 2-4 Lyon Park Road, North Ryde NSW 2113

**In a transport emergency dial 000, Police or Fire Brigade**  
**For specialist advice in an emergency only, call 1800 033 111 (24 hours)**

APVMA Approval No: 62011/20/0607  
231557  
UN-Free



**STORAGE AND DISPOSAL**

Store in the closed, original container in a dry, cool well ventilated area out of direct sunlight. Triple, or preferably pressure rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and bury empty containers in a local authority landfill. If not available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

**SAFETY DIRECTIONS**

**Harmful if inhaled or swallowed. Will irritate eyes and skin. Avoid contact with eyes and skin. DO NOT inhale spray mist. When preparing spray and using the prepared spray wear elbow-length PVC gloves and face shield. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves and face shield and contaminated clothing.**

**FIRST AID**

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone 131 126.

**MATERIAL SAFETY DATA SHEET**

If additional hazard information is required refer to the Material Safety Data Sheet. For a copy visit our website at [www.syngenta.com.au](http://www.syngenta.com.au)

**MANUFACTURER'S WARRANTY AND EXCLUSION OF LIABILITY**

Syngenta has no control over storage, handling and manner of use of this product. Where this material is not stored, handled or used correctly and in accordance with directions, no express or implied representations or warranties concerning this product (other than non-excludable statutory warranties) will apply. Syngenta accepts no liability for any loss or damage arising from incorrect storage, handling or use.

® Products marked ®, the INNOVA logo, the SYNGENTA logo and the CP FRAME are registered trademarks of a Syngenta Group Company



APVMA Approval No: 62011/20/0607  
231557

|                     |  |
|---------------------|--|
| Batch Number        |  |
| Date of Manufacture |  |

**drumMUSTER LOGO**

BARCODE

# CAUTION

KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING



# Glyphosate Trimesium 480

**HERBICIDE**

ACTIVE CONSTITUENT: 480 g/L GLYPHOSATE-TRIMESIUM

|       |          |           |
|-------|----------|-----------|
| GROUP | <b>M</b> | HERBICIDE |
|-------|----------|-----------|

*A non-selective foliar applied systemic herbicide for the control of a wide range of annual and perennial weeds as per Directions for Use*


**IMPORTANT:** Read this booklet before use

## 20 LITRES

*Syngenta Crop Protection Pty Limited  
Level 1, 2-4 Lyon Park Road, North Ryde NSW 2113*

*In a transport emergency dial 000, Police or Fire Brigade  
For specialist advice in an emergency only, call 1800 033 111 (24 hours)*

APVMA Approval No: 62011/20/0607  
231557  
UN-Free



## DIRECTIONS FOR USE

ALL STATES (except for Sugarcane - Qld, NSW, WA only)

### Restraint

DO NOT apply if heavy rain (eg 4mm) is expected within 6 hours

| Situation   | Critical Comments   |
|---|---|
| <p><b>General Weed Control</b><br/>For general weed control in and around home gardens, industrial, commercial and public service areas, parks and gardens, around farm buildings and other farm situations</p> | <p>For control of a wide range of grass and broadleaf weeds.<br/><b>Rate: 10 mL per litre of water.</b><br/>Ensure thorough coverage of weed foliage. DO NOT allow spray to contact foliage or green bark of desirable plants.<br/>Apply under good growing conditions only to actively growing weeds.</p>  |
| <p><b>Non-Cultivated Land</b><br/>Industrial, commercial, domestic and public service areas</p>   | <p>For residual control of annual weeds INNOVA GLYPHOSATE TRIMESIUM can be tank mixed with Gesaprim<sup>®</sup> Granules, Diurex* WG or Gesatop<sup>®</sup> Granules.<br/>See Compatibility Section.</p>  |
| <p><b>Forests</b><br/><b>Orchards</b><br/><b>Vines</b><br/><b>Plantations</b><br/><b>Nurseries</b></p>  | <p>INNOVA GLYPHOSATE TRIMESIUM can be used prior to the establishment of the trees and vines and amongst established trees and vines using a directed or shielded spray or wiper equipment.<br/>DO NOT allow spray or spray drift to contact green bark or stems, canes, laterals, suckers, pruned surfaces or fruit. Particular care is required to avoid spray contact with the trunks of avocado, banana, guava, kiwi fruit, lychee, mango, paw paw and stone fruit trees and ornamentals in the <i>Rosaceae</i> family.<br/>For residual control of annual weeds in citrus, pome fruit and vines INNOVA GLYPHOSATE TRIMESIUM may be tank mixed with Gesatop Granules. See Compatibility Section.</p>                    |
| <p><b>Crops</b><br/>Sown/Transplanted</p>   | <p>INNOVA GLYPHOSATE TRIMESIUM can be used to control annual and perennial weeds prior to sowing most crops. In some instances crops sown or transplanted after the use of glyphosate salts have been injured by one or more factors including inadequate soil phosphorous sorption capacity especially under dry conditions, an interaction with soil fungi and/or decaying weed residues or root exudates from treated weeds.<br/>INNOVA GLYPHOSATE TRIMESIUM applied with selective application equipment can be used to control weeds growing between crop rows and weeds extending at least 15 cm above the crop.<br/>Avoid contact of crop foliage with INNOVA GLYPHOSATE TRIMESIUM or its application equipment.</p> |
| <p><b>Sugarcane Ratoon Sprayout</b><br/>For control of sugarcane <i>Saccharum officinale</i> ratoon regrowth Qld, NSW, WA only</p>  | <p>Apply under good growing conditions only to actively growing ratoons 60 to 120 cm tall. DO NOT apply if plants are under stress from low moisture or waterlogging.<br/><b>RATE: 4 to 9 L/ha.</b> Use the lower rate for suppression or where cultivation is to follow. Use the higher rate for control without cultivation. Varieties differ in their sensitivity to INNOVA GLYPHOSATE TRIMESIUM and this can be influenced by the growing conditions.</p>   |
| <p><b>Pastures</b><br/>Spot Application<br/>Wick Wiper Application</p>  | <p>INNOVA GLYPHOSATE TRIMESIUM is non-selective and may kill desirable plants in the sprayed area.<br/>See General Instructions - Wiper Equipment.</p>  |
| <p><b>Dry drains and channels, dry margins of dams, lakes and streams</b></p>   | <p>DO NOT apply to weeds growing in or over water. DO NOT spray across open bodies of water, and DO NOT allow spray to enter the water. DO NOT allow water to return to dry channels and drains within 4 days of application.</p>   |

**Annual Weeds Controlled**

| Weed   | Rate  |
|--|---|
| <p><b>Broadleaf</b><br/>                     Bathurst Burr <i>Xanthium spinosum</i><br/>                     Bellvine <i>Ipomoea plebeia</i><br/>                     Black Bindweed <i>Polygonum convolvulus</i><br/>                     Blue Billygoat Weed <i>Ageratum houstonianum</i><br/>                     Caltrop <i>Tribulus terrestris</i><br/>                     Capeweed <i>Arctotheca calendula</i><br/>                     Carrot <i>Daucus carota</i><br/>                     Common Sowthistle <i>Sonchus oleraceus</i><br/>                     Common Thornapple <i>Datura stramonium</i><br/>                     Deadnettle <i>Lamium amplexicaule</i><br/>                     Fat Hen <i>Chenopodium album</i><br/>                     Flaxleaf Fleabane <i>Conyza bonariensis</i><br/>                     Fumitory <i>Fumaria</i> spp<br/>                     Mintweed <i>Salvia reflexa</i><br/>                     Paterson's Curse <i>Echium plantagineum</i><br/>                     Prickly Lettuce <i>Lactuca serriola</i><br/>                     Redroot Amaranth <i>Amaranthus retroflexus</i><br/>                     Rough Poppy <i>Papaver hybridum</i><br/>                     Rough Sowthistle <i>Sonchus asper</i><br/>                     Saffron Thistle <i>Carthamus lanatus</i><br/>                     Shepherd's Purse <i>Capsella bursa-pastoris</i><br/>                     Slender Celery <i>Apium leptophyllum</i><br/>                     Smooth Catsear <i>Hypochoeris glabra</i><br/>                     Spear Thistle <i>Cirsium vulgare</i><br/>                     Speedy Weed <i>Flaveria australasica</i><br/>                     Storksbill (Qld, NSW, Vic, SA, WA only) <i>Erodium</i> spp<br/>                     Tall Fleabane <i>Conyza albida</i><br/>                     Turnip Weed <i>Rapistrum rugosum</i><br/>                     Variegated Thistle <i>Silybum marianum</i><br/>                     Wild Turnip <i>Brassica tournefortii</i><br/>                     Wireweed <i>Polygonum aviculare</i> (apply to young plants<br/>                     - repeat application may be required)</p> <p><b>Grasses</b><br/>                     Annual ryegrass <i>Lolium rigidum</i><br/>                     Awnless Barnyard Grass <i>Echinochloa colona</i><br/>                     Barley Grass <i>Hordeum leporinum</i><br/>                     Barnyard Grass <i>Echinochloa crus-galli</i><br/>                     Crowsfoot Grass <i>Eleusine indica</i><br/>                     Dinebra <i>Dinebra retroflexa</i><br/>                     Grain Sorghum <i>Sorghum bicolor</i><br/>                     Great Brome <i>Bromus diandrus</i><br/>                     Lesser Canary Grass <i>Phalaris minor</i><br/>                     Liverseed Grass <i>Urochloa panicoides</i><br/>                     Maize <i>Zea mays</i><br/>                     Paradoxa Grass <i>Phalaris paradoxa</i><br/>                     Pigeon Grass <i>Setaria</i> spp<br/>                     Silver Grass <i>Vulpia</i> spp<br/>                     Soft Brome <i>Bromus mollis</i><br/>                     Sterile Brome <i>Bromus sterilis</i><br/>                     Stink Grass <i>Eragrostis cilianensis</i><br/>                     Wheat <i>Triticum aestivum</i><br/>                     Whorled Pigeon Grass <i>Setaria verticillata</i><br/>                     Wild Oats <i>Avena fatua</i>, <i>A. ludoviciana</i><br/>                     Windmill Grass <i>Chloris truncata</i><br/>                     Winter Grass <i>Poa annua</i></p> | <p><b>Boom:</b> ^2 to 3 L/ha.</p> <p><b>Handspray (handgun or knapsack):</b> 500 to 700 mL/100 L.</p> <p><b>Wiper:</b> 1 part INNOVA GLYPHOSATE TRIMESIUM to 2 parts water.</p> <p><b>^ NOTE:</b> Use the lower rate on actively growing weeds up to 15 cm high, use the higher rate on weeds which are not stressed greater than 15 cm high or when conditions are dry or cold with ongoing overcast conditions.</p> |

**Perennial Weeds Controlled**

| Weed   | State                           | Rate      |                      |                   | Critical Comments   |
|--|---------------------------------|-----------|----------------------|-------------------|---|
|  |                                 | Boom L/ha | Hand Spray per 100 L | Hand Spray per 5L |   |
| <b>Broadleaf, Brush &amp; Woody Weeds, Rushes/Sedges</b> |                                 |           |                      |                   |   |
| Boxthorn, African<br><i>Lycium ferocissimum</i>          | All States                      | -         | 1 L                  | 50 mL             | Spray foliage thoroughly. Retreatment and/or subsequent control of seedlings may be required.   |
| Bracken<br><i>Pteridium esculentum</i>                   |                                 | 9         | 1.5 L                | 75 mL             | Add Pulse* 200 mL/100 L for boom and spot spray, not wiper. Slash bracken 6 months prior to application in autumn to fully expanded, actively growing fronds. Wiper application preferred for control in pasture, refer to Wiper Equipment under General Instructions.  |
| Catsear<br><i>Hypochoeris radicata</i>                   |                                 | 3         | 700 mL               | 35 mL             | Apply at early flowering and when growing actively.   |
| Crofton Weed<br><i>Eupatorium adenophorum</i>            | Qld, NSW only                   | -         | 500 mL               | 25 mL             | Spray foliage thoroughly. Follow up control of seedlings may be required.   |
| Fennel (Aniseed)<br><i>Foeniculum vulgare</i>            | All States                      |           |                      |                   | Apply when growing actively.  |
| Lamb's Tongue<br><i>Plantago lanceolata</i>              |                                 | 3         | 700 mL               | 35 mL             | Apply when heads present and plants growing well.   |
| Nutgrass<br><i>Cyperus rotundus</i>                      |                                 | 6         | 1 L                  | 50 mL             | Apply when plants flowering in summer and growing well. Retreatment necessary for longer term control.  |
| Redflower Mallow<br><i>Modiola caroliniana</i>           |                                 | 3         | 700 mL               | 35 mL             | Apply when growing well at early flowering.   |
| Rush, Common<br><i>Juncus usitatus</i>                   |                                 |           |                      |                   | Apply when growing well. Wiper preferred for control in pasture - refer to Wiper Equipment under General Instructions. DO NOT apply to weeds growing in or over water. DO NOT spray across open bodies of water, and DO NOT allow spray to enter the water. DO NOT allow water to return to dry channels and drains within 4 days of application. |
| Salsify<br><i>Tragopogon porrifolius</i>                 |                                 | 1.5       | 500 mL               | 25 mL             | Apply at early flowering and when growing well.   |
| Sorrel<br><i>Rumex acetosella</i>                        |                                 | 4         | 1 L                  | 50 mL             |   |
| Soursob<br><i>Oxalis pes-caprae</i>                      | NSW, ACT, Vic, Tas, SA, WA only | 1.5       | 500 mL               | 25 mL             | Apply when flowering, before natural die back occurs. Retreatment required in following season for longer term control.   |
| Wild Sage<br><i>Salvia verbenaca</i>                     | All States                      |           |                      |                   | Apply at rosette to early flowering stage.  |
| Yellow Vine<br><i>Tribulis micrococcus</i>               | Qld, NSW only                   |           |                      |                   | Apply when growing actively and flowering.  |

| Weed   | State                           | Rate                                |                      |                    | Critical Comments  |
|--|---------------------------------|-------------------------------------|----------------------|--------------------|--|
|  |                                 | Boom L/ha                           | Hand Spray per 100 L | Hand Spray per 5 L |  |
| <b>Grasses</b>   |                                 |                                     |                      |                    |  |
| Carpet Grass<br><i>Axonopus affinis</i>                      | All States                      | 3                                   | 500 mL               | 25 mL              | Apply when growing well at early flowering.  |
| Common Reed<br><i>Phragmites australis</i>                   | All States                      | 9                                   | 1.3 L                | 70 mL              | Apply at early flowering. Control may not be evident until following season. Refer to Wiper equipment under General Instructions. DO NOT apply to weeds growing in or over water. DO NOT spray across open bodies of water, and DO NOT allow spray to enter the water. DO NOT allow water to return to dry channels and drains within 4 days of application. |
| Couch<br><i>Cynodon dactylon</i>                             |                                 |                                     |                      |                    | Apply when growing well at early flowering.  |
| Cumbungi<br><i>Typha domingensis</i><br><i>T. orientalis</i> |                                 | Qld, NSW, ACT, Vic, SA, WA, NT only | 6 to 9               |                    |  |
| Guinea Grass<br><i>Panicum maximum</i>                       | All States                      | 9                                   | 1.3 L                | 70 mL              | Apply when growing well at early flowering.  |
| Hamil Grass<br><i>Panicum maximum cv Hamil</i>               | All States                      | 6 to 9                              | 1 to 1.3 L           | 50 to 70 mL        | Apply when growing well at early flowering. Use the lower rate under good growing conditions. Use the higher rate for more advanced growth or under less optimum growing conditions.   |
| Humidicola<br><i>Brachiaria humidicola</i>                   |                                 |                                     |                      |                    |  |
| Johnson Grass<br><i>Sorghum halepense</i>                    |                                 | 6                                   | 1 L                  | 50 mL              | Apply when growing well at early flowering. Refer to Wiper equipment under General Instructions  |
| Kikuyu<br><i>Pennisetum clandestinum</i>                     |                                 |                                     |                      |                    | Apply when growing well at early flowering. Refer to Wiper equipment under General Instructions.   |
| Para Grass<br><i>Bracharia mutica</i>                        |                                 | 6 to 9                              | 1 to 1.3 L           | 50 to 70 mL        | Apply when growing well at early flowering. Refer to Wiper equipment under General Instructions. DO NOT apply to weeds growing in or over water. DO NOT spray across open bodies of water, and DO NOT allow spray to enter the water. DO NOT allow water to return to dry channels and drains within 4 days of application.                                  |
| Paspalum<br><i>Paspalum dilatatum</i>                        |                                 | 3                                   | 700 mL               | 35 mL              | Apply when growing well at early flowering.  |
| Phalaris<br><i>Phalaris aquatica</i>                         |                                 | NSW, ACT, Vic, SA only              | 3 to 6               | 500 mL to 1L       | 25 to 50 mL  |
| Prairie Grass<br><i>Bromus unioloides</i>                    | All States                      | 6                                   | 1 L                  | 50 mL              |  |
| Rhodes Grass<br><i>Chloris gayana</i>                        |                                 |                                     |                      |                    |  |
| Timothy Grass<br><i>Phleum pratense</i>                      | NSW, ACT, Vic, Tas, SA, WA only | 3                                   | 700 mL               | 35 mL              |  |
| Yorkshire Fog Grass<br><i>Holcus lanatus</i>                 | All States                      |                                     |                      |                    |  |

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION**

**WITHHOLDING PERIOD**

NOT REQUIRED WHEN USED AS DIRECTED

**GENERAL INSTRUCTIONS**

INNOVA GLYPHOSATE TRIMESIUM is a non-selective foliar applied herbicide which is absorbed by green plant tissue and translocated through the plant. Weeds must be growing well for optimum performance. Any conditions eg cold or overcast, which slow plant growth will delay or inhibit weed control. Avoid treating plants that are stressed from soil moisture deficit, frost, pests and diseases and water logging. Plants covered with dust may be less susceptible. INNOVA GLYPHOSATE TRIMESIUM causes yellowing and wilting of above ground parts by browning and death. Visible symptoms may take from several days to 3 to 4 weeks to appear, depending on the weed species and the weather conditions. In perennials INNOVA GLYPHOSATE TRIMESIUM controls underground roots and buds and symptoms may not be visible for a period of up to 6 weeks. Control of regrowth may not be evident until the following season.

INNOVA GLYPHOSATE TRIMESIUM is absorbed and inactivated by soil particles and will not provide residual control of weeds. Where longer term weed control is required, INNOVA GLYPHOSATE TRIMESIUM can be tank mixed with certain residual herbicides (see compatibility) or follow up applications can be made.

INNOVA GLYPHOSATE TRIMESIUM is water soluble and can be washed off plants by rain. DO NOT spray if rain is likely to occur within 6 hours. Retreatment may be necessary if heavy rain (eg 4 mm) occurs within that time.

**Mixing**

Use only clean, soft water. Use of dam water containing soil particles, or hard water eg from bores containing calcium salts may reduce efficacy.

Fill the clean spray tank 2/3 full with water. Add the required volume of INNOVA GLYPHOSATE TRIMESIUM and agitate. Complete filling the tank with the hose end under the surface to minimise frothing.

**Application**

Avoid direct or indirect contact (through spray drift) of desirable plants.

**Aerial Application**

Use the recommended rate of INNOVA GLYPHOSATE TRIMESIUM in 20 to 30 litres of water per hectare.

**Boom Equipment**

Use the recommended rate of INNOVA GLYPHOSATE TRIMESIUM in 30 to 250 litres of water per hectare. The amount of water required will vary depending on the density and size of the weeds to be controlled. Where the lower rates of INNOVA GLYPHOSATE TRIMESIUM are used for fallow weed control, use the lower water volumes. A calibrated boom fitted with flat fan nozzles operating at the optimum pressure and height for that nozzle is recommended.

**Knapsack and High Volume Equipment**

Select or adjust nozzle(s) to give a fine spray without causing undesirable runoff.

**Wiper Equipment**

INNOVA GLYPHOSATE TRIMESIUM can be applied through hand held or machine mounted wiper equipment in non-crop areas and to selectively control weeds in crops and pastures. There must be sufficient height differential between the weed and crop/pasture to allow maximum contact of the wiper with the weeds without touching desirable vegetation.

Mix 1 part INNOVA GLYPHOSATE TRIMESIUM with 2 parts clean water.

For machine mounted wipers ground speed should be 4 to 8 km/hr. Two passes, at right angles or in the opposite direction will give better results.

**Controlled Droplet Application Equipment (CDA)**

Use the following table as a guide for achieving correct application rates using the Microfit\*Herbi or similar equipment. See Species Controlled tables for specific rates and use recommendations.

For hand held equipment a walking speed of approximately 1 m/sec (3.6 km/hr) is recommended.

| Rate of INNOVA GLYPHOSATE TRIMESIUM delivered at 1 m/sec | 3 L/ha | 6 L/ha | 9 L/ha |
|--|--------|--------|--------|
| Mixture (by volume) INNOVA GLYPHOSATE TRIMESIUM: water   | 1:3    | 1:1    | 2:1    |
| Microfit*Herbi, nozzle                                   | Blue   | Blue   | Yellow |

**Compatibility**

INNOVA GLYPHOSATE TRIMESIUM can be mixed with the following herbicides and additives where recommended: Gesaprim Granules, Diurex WG or Gesatop Granules These additives may slow down initial activity of INNOVA GLYPHOSATE TRIMESIUM, but final weed control is not affected.

Addition of crystalline or liquid ammonium sulphate is not required with these products. Add the partner herbicide to the tank when 2/3 full under agitation, then the INNOVA GLYPHOSATE TRIMESIUM and complete filling. INNOVA GLYPHOSATE TRIMESIUM can be mixed with Goal\* at the spike rate of 75 mL Goal/ha to increase speed of kill and to aid control of harder-to-kill weeds such as storksbill (*Erodium* spp). Add the Goal to the 2/3 full spray tank under agitation, then the INNOVA GLYPHOSATE TRIMESIUM and complete filling. DO NOT use a tank mix of atrazine and INNOVA GLYPHOSATE TRIMESIUM for barnyard grass control.

**Additives:** Pulse penetrant: 200 mL/100 L spray solution for use with INNOVA GLYPHOSATE TRIMESIUM against bracken only.

As formulations of other manufacturers' products are beyond the control of Syngenta Crop Protection, and water quality varies with location, all mixtures should be tested prior to mixing commercial quantities.

|       |   |           |
|-------|---|-----------|
| GROUP | M | HERBICIDE |
|-------|---|-----------|

### Resistant Weeds Warning

INNOVA GLYPHOSATE TRIMESIUM 480 Herbicide is a member of the Glycines group of herbicides. INNOVA GLYPHOSATE TRIMESIUM has the inhibition of EPSP synthase mode of action. For weed resistance management INNOVA GLYPHOSATE TRIMESIUM is a Group M herbicide. Some naturally occurring weed biotypes resistant to INNOVA GLYPHOSATE TRIMESIUM and other Group M herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by INNOVA GLYPHOSATE TRIMESIUM or other Group M herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use, Syngenta Crop Protection Pty Limited accepts no liability for any losses that may result from the failure of INNOVA GLYPHOSATE TRIMESIUM to control resistant weeds.

### PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

DO NOT apply under weather conditions or from spraying equipment which may cause spray to drift onto nearby susceptible plants/crops, cropping lands or pastures.

### PROTECTION OF LIVESTOCK

Low hazard to bees. No special precautions are required.

### PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate dams, rivers or streams with the product or used container. DO NOT apply to weeds growing in or over water. DO NOT spray across open bodies of water.

### STORAGE AND DISPOSAL

Store in the closed, original container in a dry, cool well ventilated area out of direct sunlight. Triple, or preferably pressure rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and bury empty containers in a local authority landfill. If not available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

### SAFETY DIRECTIONS

**Harmful if inhaled or swallowed. Will irritate eyes and skin. Avoid contact with eyes and skin. DO NOT inhale spray mist. When preparing spray and using the prepared spray wear elbow-length PVC gloves and face shield. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves and face shield and contaminated clothing.**

### FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone 131 126.

### MATERIAL SAFETY DATA SHEET

If additional hazard information is required refer to the Material Safety Data Sheet. For a copy visit our website at [www.syngenta.com.au](http://www.syngenta.com.au)

**MANUFACTURER'S WARRANTY AND EXCLUSION OF LIABILITY**

Syngenta has no control over storage, handling and manner of use of this product. Where this material is not stored, handled or used correctly and in accordance with directions, no express or implied representations or warranties concerning this product (other than non-excludable statutory warranties) will apply. Syngenta accepts no liability for any loss or damage arising from incorrect storage, handling or use.

® Products marked ®, the INNOVA logo,  
the SYNGENTA logo and the CP FRAME  
are registered trademarks of a Syngenta Group  
Company



\* Registered trademark