

DIRECTIONS FOR USE

Restraint

DO NOT treat physically damaged seed.

Crop	Disease/Pest	Rate	Critical Comments
Wheat	Bunt Flag Smut (seed and soil borne) Loose Smut	100 mL/100 kg seed	INNOVA TEBUCONAZOLE 25C should be mixed with water to give even coverage of seed. The quantity of water used for mixing will vary depending on type of equipment and quality of seed. Use a minimum of 400 mL of mixture (ie INNOVA TEBUCONAZOLE 25C plus water) with each 100 kg of seed. Do not use more than 600 mL of mixture (product plus water) with each 100 kg of seed. Whatever dilution is used it is essential that 100 mL of INNOVA TEBUCONAZOLE 25C is applied per 100 kg seed. The mixture should be gently stirred regularly. If nasal irritation occurs wear a face mask while treating seed.
Barley Oats	Covered and Loose Smut		
Barley Oats Wheat	Protection against pests of stored grains: Granary Weevil, Indian Meal Moth, Lesser Grain Borer, Rice Weevil, Rust-red Flour Beetle, Sawtoothed Grain Beetle, Tropical Warehouse Moth		

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION

WITHHOLDING PERIOD

NOT REQUIRED WHEN USED AS DIRECTED

GENERAL INSTRUCTIONS

INNOVA TEBUCONAZOLE 25C may delay emergence of wheat. Crops will recover rapidly in most soils without any effect on yield. Sowing INNOVA TEBUCONAZOLE 25C treated wheat deeper than 50 mm may reduce plant establishment in heavier soils or soils that will crust or clod at the surface after sowing. Sowing rates may need to be adjusted accordingly.

Seed Quality

INNOVA TEBUCONAZOLE 25C should not be used on seed with more than 12% moisture content, or on sprung, sprouted, damaged or severely pinched seed, or seed of poor viability. If in doubt, have a germination test carried out on the seed before treatment to ensure that it is of acceptable standard.

Treating with INNOVA TEBUCONAZOLE 25C as directed will raise the moisture level of the seed by up to 0.6%, depending on weather conditions at treatment. The use of INNOVA TEBUCONAZOLE 25C mixed with water at recommended rates will have no effect on the storage life of treated sound seed.

Storage of Treated Seed

Treated seed may be held over for sowing in the following season. If so, additional insecticide treatment may be necessary. If the seed is not used immediately after treatment it should be stored in a dry, well-ventilated place. Although Innova Tebuconazole C has no effect on the viability of treated seed, subsequent germination can be adversely affected by poor storage conditions such as high moisture combined with high temperatures. No liability can therefore be accepted for the performance of stored treated seed.

Cleaning Up

Equipment should be thoroughly cleaned with water. In the event of changing from this water based flowable formulation to a solvent-based liquid formulation, it is essential to rinse equipment with methylated spirits after thoroughly cleaning with water. If changing from a solvent-based liquid to this water based flowable thoroughly clean with methylated spirits then rinse with water. Failure to do so will result in clogging of equipment.

GROUP	C	FUNGICIDE
--------------	----------	------------------

Fungicide Resistance Warning

INNOVA TEBUCONAZOLE 25C Flowable Seed Dressing is a member of the DMI group of fungicides. For fungicide resistance management the product is a Group C fungicide. Some naturally occurring individual fungi resistant to the product and other Group C fungicides may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungal population if these fungicides are used repeatedly. These resistant fungi will not be controlled by this product and other Group C fungicides, thus resulting in a reduction in efficacy and possible yield loss. Since the occurrence of resistant fungi is difficult to detect prior to use, Syngenta Crop Protection Pty Limited accepts no liability for any losses that result from failure of this product to control resistant fungi.

Insecticide Resistance Warning

GROUP 3A INSECTICIDE

For insecticide resistance management INNOVA TEBUCONAZOLE 25C Flowable Seed Dressing is a Group 3A insecticide. Some naturally occurring insect biotypes resistant to INNOVA TEBUCONAZOLE 25C and other Group 3A insecticides may exist through normal genetic variability in any insect population. The resistant individuals can eventually dominate the insect population if INNOVA TEBUCONAZOLE 25C or other Group 3A insecticides are used repeatedly. The effectiveness of INNOVA TEBUCONAZOLE 25C on resistant individuals could be significantly reduced. Since occurrence of resistant individuals is difficult to detect prior to use, Syngenta Crop Protection Pty Limited accepts no liability for any losses that may result from the failure of INNOVA TEBUCONAZOLE 25C to control resistant insects. INNOVA TEBUCONAZOLE may be subject to specific resistance management strategies. For further information contact your local supplier or local agricultural department agronomist.

Precaution

DO NOT allow seed treated with this product to contaminate seed intended for human consumption. If possible, use a separate auger for treating seed. If using the same auger for treating seed and moving grain for human consumption, remove all INNOVA TEBUCONAZOLE residue from the auger to avoid contaminating untreated seed with dressing residues.

PROTECTION OF LIVESTOCK

Seed treated with this product must not be used for animal consumption or poultry feed or mixed with animal feed. Do NOT allow seed treated with this product to contaminate seed intended for animal consumption.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT feed treated seed or otherwise expose to wild or domestic birds. Harmful to fish. DO NOT contaminate dams, rivers, ponds, waterways or drains with this product, used containers or bags which have held treated seed.

STORAGE AND DISPOSAL

Product and original container

Store product in the closed original container in a dry, cool, well ventilated area out of direct sunlight. Single rinse before disposal. Add rinsings to disposal pit. Do not dispose of undiluted chemicals on-site. Puncture and bury empty containers in a local authority landfill. If not available, bury containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and roots. Empty containers and product should not be burnt.

Treated seed and containers of treated seed

When treated seed is stored it should be kept apart from other grain and the bags or other containers should be clearly marked to indicate the contents have been treated with this product. DO NOT use treated seed for human consumption. Bags which have held treated seed are not be used for any other purpose.

SAFETY DIRECTIONS

May irritate the eyes. Avoid contact with eyes. Wash hands after use.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone 131 126.

MATERIAL SAFETY DATA SHEET

If additional hazard information is required refer to the Material Safety Data Sheet. For a copy visit our website at www.syngenta.com.au

MANUFACTURER'S WARRANTY AND EXCLUSION OF LIABILITY

Syngenta has no control over storage, handling and manner of use of this product. Where this material is not stored, handled or used correctly and in accordance with directions, no express or implied representations or warranties concerning this product (other than non-excludable statutory warranties) will apply. Syngenta accepts no liability for any loss or damage arising from incorrect storage, handling or use.

® The INNOVA logo, the SYNGENTA logo and the CP FRAME are registered trademarks of a Syngenta Group Company
APVMA Approval No: 62155/10/0707
ITC0507



Batch No	
Date of Manufacture	