

## READ SAFETY DIRECTIONS BEFORE OPENING OR USING



### POLYMER-BINDER SEED COATING

Adjuvant to enhance the appearance, improve seed flow and prevent the loss of seed treatments during packing, transportation and drilling of seeds

Pack size: 200 L

#### GENERAL INSTRUCTIONS

NACRET is a polymer-binder for film coating use during the application process of most insecticide and fungicide seed treatments. NACRET will enhance the appearance, improve the seed flow in the drill, and prevent loss of seed treatment products during packaging, transportation and drilling of treated seed and will result in improved operator safety.

##### APPLICATION EQUIPMENT

NACRET can be applied in standard continuous flow equipment without special modifications.

It is essential that the continuous flow equipment is calibrated before commercial use, as flow characteristics may not be the same as for other seed treatments.

##### APPLICATION EQUIPMENT CLEANUP

Thoroughly clean all equipment after use of this product with hot water.

##### COMPATIBILITY

NACRET diluted in water can be mixed with other seed dressing products directly in the slurry tank.

##### PROTECTION OF LIVESTOCK

DO NOT feed treated seed to animals, including poultry.  
DO NOT allow treated seed to contaminate grain intended for animal consumption.

##### PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate streams, rivers or waterways with the product or used containers.

##### STORAGE AND DISPOSAL

When treated seed is stored it should be kept apart from other grain and the bags or other containers should be clearly marked to indicate the contents have been treated with NACRET. Bags which have held treated seed should not be used for any other purpose.

Store in the closed, original container in a cool, well ventilated area, keep as cool as possible.

DO NOT store for prolonged periods in direct sunlight.  
DO NOT use treated seed for human consumption.  
DO NOT allow treated seed to contaminate seed intended for human consumption.  
DO NOT use treated seed for the production of oil.

Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point.

If not recycling, break, crush, or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

#### SAFETY DIRECTIONS

Avoid contact with eyes and skin. DO NOT inhale spray mist. When using the product and while handling treated seeds wear cotton overalls buttoned to the neck and wrist, washable hat and impervious footwear. If product gets on skin immediately wash area with soap and water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use wash contaminated clothing.

#### FIRST AID

If poisoning occurs contact a doctor or Poisons Information Centre. Phone 131 126.

#### MANUFACTURER'S WARRANTY AND EXCLUSION OF LIABILITY

Syngenta has no control over storage, handling and manner of use of this product. Where this material is not stored, handled or used correctly and in accordance with directions, no express or implied representations or warranties concerning this product (other than non-excludable statutory warranties) will apply. Syngenta accepts no liability for any loss or damage arising from incorrect storage, handling or use.

TM Trademark of LA QUINOLEINE, Paris, France



## DIRECTIONS FOR USE

Crop	Rate per 100 kg seed mL NACRET	Critical Comments
Field Peas, Soybeans, Beans	200 to 300	<ol style="list-style-type: none"><li>1. Pour the recommended quantity of water in the spray tank and add the required amount of the seed treatment as per its Directions for Use table.</li><li>2. Add the required amount of NACRET while stirring the solution.</li><li>3. The solution is ready for use after a few minutes when homogeneous.</li><li>4. Apply the prepared solution to the seed to be treated as per the instructions given on the seed treatment label.</li></ol>
Sunflowers	300 to 750	
Canola	750 to 1000	
Maize	250 to 1000	
Vegetable seeds	1000 to 3000	
Cotton	300 to 500	

**WITHHOLDING PERIOD: NOT REQUIRED WHEN USED AS DIRECTED.**