

POISON

KEEP OUT OF REACH OF CHILDREN

READ SAFETY DIRECTIONS BEFORE OPENING OR USING



Proclaim[®]
Insecticide

syngenta.

**ACTIVE CONSTITUENT: 44 g/kg EMAMECTIN present as
EMAMECTIN BENZOATE**

GROUP 6 INSECTICIDE

*For the control of Diamondback Moth (Cabbage Moth) and
Cabbage White Butterfly in Brassica Vegetables, Light Brown
Apple Moth and Grapevine Moth in Grapes and Heliothis in
Capsicum, Lettuce, Sweet Corn and Tomatoes*

Syngenta Crop Protection Pty Limited
Level 1, 2-4 Lyonpark Road, Macquarie Park, NSW 2113

**In a transport emergency dial 000, Police or Fire Brigade
For specialist advice in an emergency only, call 1800 033 111 (24 hours)**

APVMA Approval No: 50919/600 or 3/1008

TM

DIRECTIONS FOR USE

Crop	Pest	Rate	Critical Comments
Brassica Vegetables Broccoli, Brussels Sprout, Cabbages, Cauliflower only	Diamondback Moth (Cabbage Moth) (<i>Plutella xylostella</i>) and Cabbage White Butterfly (<i>Pieris rapae</i>)	250 to 300 g/ha	Spray at first signs of insect infestation as indicated by local spray thresholds. Use the lower rate on low to moderate infestations. Add Citowett* or Agral® at the appropriate label rate. DO NOT make more than 4 applications to any brassica crop. Where more than one crop is grown DO NOT make more than 4 applications of PROCLAIM in any one year. PROCLAIM should be used according to CropLife insecticide resistance strategies.
Capsicum, Lettuce	Heliothis (<i>Helicoverpa armigera</i> and <i>Helicoverpa punctigera</i>)	150 to 250 g/ha	Spray at first signs of insect infestation as indicated by local spray thresholds. For best results apply soon after the <i>Helicoverpa</i> eggs have hatched. DO NOT apply more than 4 sprays of Proclaim per crop. Where more than one crop is grown per year DO NOT apply more than 4 sprays per year. Ensure thorough spray coverage. Use the lower rate when targeting light infestations of small larvae. Use 250 g/ha during periods of heavy insect pressure or under very hot and sunny conditions. PROCLAIM should be used according to the CropLife resistance strategies.
Grapes Except grapes grown for dried fruit production	Lightbrown Apple Moth, Grapevine Moth	Dilute Spraying 15 g/100 L Concentrate Spraying Refer to the Mixing/ Application Section	DO NOT apply after bunch closure. For optimal control of Light Brown Apple Moth apply soon after egg lay when larvae are small in size and before larvae become webbed into the bunches. Eggs laid after application on new growth may not be controlled. For Grapevine Moth control, spray when local threshold levels have been reached. Monitor pest levels after application, a further application may be required (minimum re-treatment interval: 7 to 14 days). Apply by dilute or concentrate spraying equipment. Apply a maximum of 2 applications. Ensure thorough coverage. Apply the same amount of product to the target crop whether applying this product by dilute or concentrate spraying methods.
Sweet Corn	Heliothis (<i>Helicoverpa armigera</i> , <i>Helicoverpa punctigera</i>)	150 to 250 g/ha	For best results apply just prior to or just after the <i>Helicoverpa</i> eggs have hatched, but before larvae enter the cob. DO NOT apply more than 4 sprays of PROCLAIM per crop. Where more than one crop is grown per year DO NOT apply more than 4 sprays per year. Thorough spray coverage is critical. Larvae present within the cob at the time of spraying may not be controlled. Use the lower rate when targeting light infestations of small larvae. PROCLAIM should be used according to the CropLife resistance strategies.

Crop	Pest	Rate	Critical Comments
Tomatoes	Heliothis (<i>Helicoverpa armigera</i> , <i>Helicoverpa punctigera</i>)	150 to 250 g/ha or 15 to 25 g/100 L (dilute spraying) for trellised Tomatoes. Concentrate Spraying Refer to the Mixing/ Application section	Spray at first signs of insect infestation as indicated by local spray thresholds. For best results apply soon after the <i>Helicoverpa</i> eggs have hatched. DO NOT apply more than 4 sprays of PROCLAIM per crop. Where more than one crop is grown per year DO NOT apply more than 4 sprays per year. Ensure thorough spray coverage. Use the lower rate when targeting light infestations of small larvae. Use 250 g/ha during periods of heavy insect pressure or under very hot and sunny conditions. Trellised Tomatoes: In the case of dilute spraying (g/100 L) apply in the range of 400 to 500 L/ha after transplanting and increase to 800 to 1000 L/ha at full canopy. PROCLAIM should be used according to the CropLife resistance strategies.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION

WITHHOLDING PERIODS

Brassica Vegetables: DO NOT HARVEST FOR 3 DAYS AFTER APPLICATION
DO NOT USE TREATED CROP, CROP WASTE OR PRODUCE FOR STOCK FOOD

Capsicum, Lettuce, Tomatoes:
DO NOT HARVEST FOR 3 DAYS AFTER APPLICATION
DO NOT ALLOW LIVESTOCK TO GRAZE TREATED CROPS OR CROP STUBBLE

Grapes:
DO NOT HARVEST FOR 8 WEEKS AFTER APPLICATION
DO NOT GRAZE TREATED AREAS, OR CUT FOR STOCKFEED, FOR 8 WEEKS AFTER APPLICATION

Sweet Corn:
DO NOT HARVEST FOR 3 DAYS AFTER APPLICATION
DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 21 DAYS AFTER APPLICATION

EXPORT TRADE ADVICE

Treated crop commodities destined for export may require extra time being allowed between application and harvest, as some export market have either no Maximum Residue Limits (MRLs) or different MRLs to those of Australia. Details of overseas standards and any export interval can be obtained by contacting Syngenta Crop Protection before using this product.

GENERAL INSTRUCTIONS

Mixing

Add the required amount of PROCLAIM to a partly filled spray tank, and then add the remainder of the water.

Application

PROCLAIM is not systemic making good coverage essential - ensure thorough coverage of the crop
DO NOT apply by aerial application

Brassica Vegetables: Apply PROCLAIM in a minimum of 400 L water/ha

Capsicums, Lettuce, Sweet Corn, Tomatoes: Ensure thorough coverage of the crop

Grapes: Apply PROCLAIM in a minimum of 250 L water/ha

Mixing/Application – Grapes and Trellised Tomatoes

Dilute Spraying: Use a sprayer designed to apply high volumes of water up to the point of runoff and matched to the crop being sprayed. Set up and operate the sprayer to achieve even coverage throughout the crop canopy. Apply sufficient water to cover the crop to the point of runoff. Avoid excessive runoff. The required water volume may be determined by applying different test volumes, using different settings on the sprayer, from industry guidelines or expert advice. Add the amount of product specified in the Directions for Use table for each 100 L of water. Spray to the point of runoff. The required dilute spray volume will change and the sprayer set up and operation may also need to be changed as the crop grows.

Concentrate Spraying: Use a sprayer designed and set up for concentrate spraying (that is a sprayer which applies water volumes less than those required to reach the point of runoff) and matched to the crop being sprayed. Set up and operate the sprayer to achieve even coverage throughout the crop canopy using your chosen water volume. Determine an appropriate dilute spray volume (see Dilute Spraying) for the crop canopy. This is needed to calculate the concentrate mixing rate.

The mixing rate for concentrate spraying can then be calculated in the following way:

Example Only

1. Dilute spray volume as determined above: For example 1000 L/ha
2. Your chosen concentrate spray volume: For example 500 L/ha
3. The concentration factor in this example is: 2 X (ie. $1000 \text{ L} \div 500 \text{ L} = 2$)
4. If the dilute label rate is 15 g/100 L, then the concentrate rate becomes 2 x 15, that is 30 g/100 L of concentrate spray

The chosen spray volume, amount of product per 100 L of water, and the sprayer set up and operation may need to be changed as the crop grows.

For further information on concentrate spraying, users are advised to consult relevant industry guidelines, undertake appropriate competency training and follow industry Best Practices.

Compatibility

PROCLAIM is compatible with Chess[®] Insecticide, Delfin^{*} WG Insecticide, Pirimor[®] WG Aphicide, Supracide[®] 400 EC, Topas[®] 100 EC Systemic Fungicide, Citowett, Agral, Confidor^{*} 200SC Insecticide, Mavrik^{*} Aquaflo Insecticide and mancozeb. As formulations of other manufacturer's products are beyond the control of Syngenta, and the quality of water may vary with location, all mixtures should be tested prior to mixing commercial quantities.

Insecticide Resistance Warning

GROUP	6	INSECTICIDE
-------	---	-------------

For insecticide resistance management PROCLAIM Insecticide is a Group 6 insecticide. Some naturally occurring insect biotypes resistant to PROCLAIM and other Group 6 insecticides may exist through normal genetic variability in any insect population. The resistant individuals can eventually dominate the insect population if PROCLAIM and other Group 6 insecticides are used repeatedly. The effectiveness of PROCLAIM on resistant individuals could be significantly reduced. Since occurrence of resistant individuals is difficult to detect prior to use, Syngenta Crop Protection accepts no liability for any losses that may result from failure of PROCLAIM to control resistant insects. PROCLAIM may be subject to specific resistance management strategies. For further information contact your supplier, Syngenta Crop Protection representative or local agricultural department agronomist.

PRECAUTION

Re-entry Period: DO NOT allow entry into treated areas for 8 hours or until the spray has dried. If prior entry is necessary, wear chemical resistant gloves.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

Dangerous to fish and other aquatic organisms. DO NOT contaminate streams, rivers or waterways with the chemical or used containers. DO NOT apply under weather conditions or from spraying equipment which could be expected to cause spray to drift onto adjacent areas, particularly wetlands, water bodies or watercourses.

PROTECTION OF LIVESTOCK

Dangerous to bees. DO NOT spray any plants in flower while bees are foraging.

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well ventilated area. DO NOT store for prolonged periods in direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to the spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

SAFETY DIRECTIONS

Harmful if swallowed. Will irritate the eyes and skin. Avoid contact with eyes and skin. When opening the container and preparing the spray wear:

- cotton overalls buttoned to the neck and wrist (or equivalent clothing)
- elbow length PVC gloves
- goggles or face shield

Wash hands after use. After each day's use, wash gloves, goggles or face shield and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone 131 126.

MATERIAL SAFETY DATA SHEET

If additional hazard information is required refer to the Material Safety Data Sheet. For a copy phone 1800 067 108 or visit our website at www.syngenta.com.au

MANUFACTURER'S WARRANTY AND EXCLUSION OF LIABILITY

Syngenta has no control over storage, handling and manner of use of this product. Where this material is not stored, handled or used correctly and in accordance with directions, no express or implied representations or warranties concerning this product (other than non-excludable statutory warranties) will apply. Syngenta accepts no liability for any loss or damage arising from incorrect storage, handling or use.

Product names marked ® or ™, the ALLIANCE FRAME
the SYNGENTA Logo and the PURPOSE ICON
are Trademarks of a Syngenta Group Company



* Trademark (statement only to appear in booklet)